# Sensory Overload Soothing Strategies Checklist A calming guide for children and adults with autism



#### **✓** Reduce the Environment

- $\square$  Move to a quieter space
- □ Dim lights or use natural lighting
- □ Turn off loud appliances (vacuum, blender, TV)
- □ Reduce visual clutter or step away from crowds
- □ Offer sunglasses or a hat for bright lighting



## **✓** Soothing Tools & Supports

- □ Noise-canceling headphones
- □ Soft, calming music
- □ Weighted blanket, lap pad, or compression vest
- □ Fidget toy or sensory object
- □ Chewelry or oral sensory tool
- □ Soft textured comfort item (plush, textured cloth)

## ✓ Calming the Body

- □ Deep breaths together (5 slow breaths)
- ☐ Gentle deep pressure (if preferred)
- □ Hand squeezes or "palm press"
- □ Slow rocking or swaying
- □ Stretching or simple movement breaks
- □ Access to a quiet "calm space"

## √ Reassurance & Regulation

- ☐ Stay close and present
- ☐ Keep instructions short and gentle
- □ Remind them they are safe
- □ Provide a familiar object (blanket, toy, device)
- ☐ Maintain predictable routine afterward
- ☐ Allow recovery time before returning to tasks

#### ✓ After the Overload

- ☐ Ask simple reflection questions when calm
- "Was it too loud?"
- "Did that feel like too much?"
- □ Note the trigger for future planning
- □ Adjust environment or routine to prevent repeat stress
- ☐ Celebrate their effort and recovery

#### **✓** Supporting Communication

- □ Offer simple choices: "Do you want quiet or dark?"
- ☐ Use visuals or gestures instead of words
- ☐ Give extra time to respond
- ☐ Avoid asking too many questions
- □ Use a calm, predictable tone
- □ Validate feelings: "It's okay. That was a lot."